

# The Fate of the Landau Levels under Perturbations of Constant Sign

June 30, 2009

FRÉDÉRIC KLOPP, GEORGI RAIKOV

## Abstract

We show that the Landau levels cease to be eigenvalues if we perturb the 2D Schrödinger operator with constant magnetic field, by bounded electric potentials of fixed sign. We also show that, if the perturbation is not of fixed sign, then any Landau level may be an eigenvalue of the perturbed problem.

**AMS 2000 Mathematics Subject Classification:** 35J10, 81Q10, 35P20

**Keywords:** Landau Hamiltonians, splitting of Landau levels

## 1 Introduction. Main results

In this note we consider the Landau Hamiltonian  $H_0$ , i.e. the 2D Schrödinger operator with constant magnetic field. It is well-known that the spectrum of  $H_0$  consists of an arithmetic progression of eigenvalues called *Landau levels* of infinite multiplicity. In Theorem 1 we show that under perturbations by fairly general electric potentials of constant sign, the Landau levels cease to be eigenvalues of the perturbed operator. Moreover, in Theorem 2 we show that for each fixed Landau level there exist non-constant-sign electric potentials such that the Landau levels is still an eigenvalue of infinite multiplicity of the perturbed operator.

Let

$$H_0 := \left( -i \frac{\partial}{\partial x} + \frac{by}{2} \right)^2 + \left( -i \frac{\partial}{\partial y} - \frac{bx}{2} \right)^2 - b$$

be the Landau Hamiltonian shifted by the value  $b > 0$  of the constant magnetic field. The operator  $H_0$  is self-adjoint in  $L^2(\mathbb{R}^2)$ , and essentially self-adjoint on  $C_0^\infty(\mathbb{R}^2)$ . Note that  $C_0^\infty(\mathbb{R}^2) \setminus \{0\}$  is a form core for the operator  $H_0$ . It is well-known (see [3, 6, 1]) that the spectrum  $\sigma(H_0)$  of the operator  $H_0$  consists of the so-called Landau levels  $2bq$ ,  $q \in \mathbb{N} := \{0, 1, 2, \dots\}$ , which are eigenvalues of  $H_0$  of infinite multiplicity. Let  $V \in L^\infty(\mathbb{R}^2; \mathbb{R})$ . We will suppose that

$$c\chi(\mathbf{x}) \leq V(\mathbf{x}), \quad \mathbf{x} = (x, y) \in \mathbb{R}^2, \quad (1.1)$$

where  $c > 0$  is a constant and  $\chi$  is the characteristic function of a disk of radius  $r > 0$  in  $\mathbb{R}^2$ , and

$$\|V\|_{L^\infty(\mathbb{R}^2)} < 2b. \quad (1.2)$$

Set  $H_\pm := H_0 \pm V$ . The main result of the note is the following

**Theorem 1.** *Fix  $q \in \mathbb{N}$ .*

(i) *Assume that  $V \in L^\infty(\mathbb{R}^2; \mathbb{R})$  satisfies (1.1); if  $q \geq 1$ , suppose in addition that (1.2) holds true. Then we have*

$$\text{Ker}(H_+ - 2bq) = \{0\}. \quad (1.3)$$

(ii) *Assume that  $V$  satisfies (1.1) and (1.2). Then we have*

$$\text{Ker}(H_- - 2bq) = \{0\}. \quad (1.4)$$

The proof of Theorem 1 can be found in Section 2.

To the authors' best knowledge the fate of the Landau levels under perturbations of the described class had never been addressed in the mathematical literature. However, the asymptotic distribution of the discrete spectrum near the Landau levels of various perturbations of the Landau Hamiltonian and its generalizations has been considered by numerous authors (see [11, 4, 12, 7, 2, 10, 15, 14, 9]); in particular, it was shown in [12] that for any  $V$  which satisfies (1.1), and is relatively compact with respect to  $H_0$ , and for any Landau level there exists an infinite sequence of discrete eigenvalues of  $H_\pm$  which accumulates to this Landau level. Such results are related to the problem treated here: indeed, the existence of such an infinite sequence is a necessary condition that the Landau level not be an infinitely degenerate eigenvalue of  $H_\pm$ .

The fact that  $V$  has a fixed sign plays a crucial role in our result, as shows the following

**Theorem 2.** *Fix  $q \geq 0$ . Then, there exists a bounded compactly supported non-constant-sign potential  $V$  such that  $\|V\|_{L^\infty(\mathbb{R}^2)} < b$  and*

$$\dim \text{Ker}(H_0 + V - 2bq) = \infty. \quad (1.5)$$

The proof of Theorem 2 is contained in Section 3. Its strategy is to consider radially symmetric potentials  $V$ , and, applying a decomposition into a Fourier series with respect to the angular variable, to represent the operator  $H_0 + V$  as an infinite sum of ordinary differential operators involving only the radial variable. Such a representation of  $H_0 + V$  is well known, and has been used in different contexts of the spectral theory of the perturbed Landau Hamiltonian (see e.g. [1, 8]). To prove Theorem 2, the basic consequence is that, for a compactly supported, radially symmetric potential  $V$ , the first derivative with respect to the coupling constant  $\lambda$  at  $\lambda = 0$  of the eigenvalues of  $H_0 + \lambda V$  close to the  $q$ -th Landau level is determined by  $V$  near the external rim of its support. Thus, writing  $V = V_t$  as an infinite sum of concentric potentials depending on different coupling constant  $t = (t_l)_{l \geq 1} \in \ell^\infty(\mathbb{N}^*)$ , one can construct an analytic mapping

from a neighborhood of 0 in  $\ell^\infty(\mathbb{N}^*)$  to a subset of the eigenvalues of  $H_0 + V_t$  near the  $q$ -th Landau level, the Jacobian of which we control for  $t = 0$ .

The potential exhibited in Theorem 2 can be chosen arbitrarily small. Following the same idea, one can also construct compactly supported potentials such that any of the Landau levels be of finite non trivial multiplicity or non compactly supported, bounded potentials such that (1.5) be satisfied for any  $q \in \mathbb{N}$ .

## 2 Proof of Theorem 1

Denote by  $\Pi_q$ ,  $q \in \mathbb{N}$ , the orthogonal projection onto  $\text{Ker}(H_0 - 2bq)$ . Set

$$\Pi_q^+ := \sum_{j=q}^{\infty} \Pi_j, \quad \Pi_q^- := I - \Pi_q^+, \quad q \in \mathbb{N}.$$

In order to prove Theorem 1, we need a technical result concerning some Toeplitz-type operators of the form  $\Pi_q V \Pi_q$ .

**Lemma 3.** *Let  $V \in L^\infty(\mathbb{R}^2; \mathbb{R})$  satisfy (1.1). Fix  $q \in \mathbb{N}$ . Then*

$$\langle \Pi_q V \Pi_q u, u \rangle = 0, \quad u \in L^2(\mathbb{R}^2), \quad (2.1)$$

where  $\langle \cdot, \cdot \rangle$  denotes the scalar product in  $L^2(\mathbb{R}^2)$ , implies

$$\Pi_q u = 0. \quad (2.2)$$

*Proof.* By (1.1) and (2.1),

$$0 \leq c \langle \Pi_q \chi \Pi_q u, u \rangle \leq \langle \Pi_q V \Pi_q u, u \rangle = 0, \quad (2.3)$$

i. e.

$$\langle \Pi_q \chi \Pi_q u, u \rangle = 0. \quad (2.4)$$

Denote by  $T := \Pi_q \chi \Pi_q$  the operator self-adjoint in the Hilbert space  $\Pi_q L^2(\mathbb{R}^2)$ . The operator  $T$  is positive and compact, and its eigenvalues can be calculated explicitly (see [12, Eq. (3.32)]). This explicit calculation implies that  $\text{Ker } T = \{0\}$ . Therefore, (2.2) follows from (2.4).  $\square$

*Proof of Theorem 1.* First, we prove (1.3) in the case  $q = 0$ . Assume that there exists  $u \in D(H_+) = D(H_0)$  such that  $H_+ u = 0$ . Hence,

$$\langle H_0 u, u \rangle + \langle V u, u \rangle = 0. \quad (2.5)$$

The two terms at the l.h.s. of (2.5) are non-negative, and therefore they both should be equal to zero. Since  $\langle H_0 u, u \rangle = 0$ , we have

$$u = \Pi_0 u. \quad (2.6)$$

Therefore,  $\langle Vu, u \rangle = \langle \Pi_0 V \Pi_0 u, u \rangle = 0$ . By Lemma 3, we have  $\Pi_0 u = 0$ , and by (2.6) we conclude that  $u = 0$ .

Next, we prove (1.3) in the case  $q \geq 1$ . Assume that there exists  $u \in D(H_0)$  such that

$$H_+ u = 2bqu. \quad (2.7)$$

Set  $u_+ := \Pi_q^+ u$ ,  $u_- := u - u_+$ ; evidently,  $u_{\pm} \in D(H_0)$ . Since  $H_0$  commutes with the projections  $\Pi_q^{\pm}$ , (2.7) implies

$$H_0 u_+ - 2bqu_+ + \Pi_q^+ V \Pi_q^+ u_+ + \Pi_q^+ V \Pi_q^- u_- = 0, \quad (2.8)$$

$$H_0 u_- - 2bqu_- + \Pi_q^- V \Pi_q^- u_- + \Pi_q^- V \Pi_q^+ u_+ = 0. \quad (2.9)$$

Now note that the operator  $H_0 + \Pi_q^- V \Pi_q^- - 2bq$  is boundedly invertible in  $\Pi_q^- L^2(\mathbb{R}^2)$ , and its inverse is a negative operator. Moreover, by (2.9) we have

$$u_- = - (H_0 + \Pi_q^- V \Pi_q^- - 2bq)^{-1} \Pi_q^- V \Pi_q^+ u_+, \quad (2.10)$$

which inserted into (2.8) implies

$$H_0 u_+ - 2bqu_+ + \Pi_q^+ V \Pi_q^+ u_+ - \Pi_q^+ V \Pi_q^- (H_0 + \Pi_q^- V \Pi_q^- - 2bq)^{-1} \Pi_q^- V \Pi_q^+ u_+ = 0,$$

and hence,

$$\begin{aligned} & \langle (H_0 - 2bq)u_+, u_+ \rangle + \langle \Pi_q^+ V \Pi_q^+ u_+, u_+ \rangle \\ & - \langle \Pi_q^+ V \Pi_q^- (H_0 + \Pi_q^- V \Pi_q^- - 2bq)^{-1} \Pi_q^- V \Pi_q^+ u_+, u_+ \rangle = 0. \end{aligned} \quad (2.11)$$

The three terms on the l.h.s. of (2.11) are non-negative, and hence they all should be equal to zero. Since  $u_+ = \Pi_q^+ u_+$ , the equality  $\langle (H_0 - 2bq)u_+, u_+ \rangle = 0$  implies

$$u_+ = \Pi_q u_+. \quad (2.12)$$

Therefore,  $\langle \Pi_q^+ V \Pi_q^+ u_+, u_+ \rangle = \langle \Pi_q V \Pi_q u_+, u_+ \rangle$ , and  $\langle \Pi_q^+ V \Pi_q^+ u_+, u_+ \rangle = 0$  is equivalent to  $\langle \Pi_q V \Pi_q u_+, u_+ \rangle = 0$ . Now by Lemma 3 we have  $\Pi_q u_+ = 0$ , by (2.12) we have  $u_+ = 0$ , and by (2.10) we have  $u_- = 0$ . Therefore,  $u = 0$ .

Finally, we sketch the proof of (1.4) which is quite similar to the one of (1.3). Let  $w \in D(H_0)$ ,  $H_- w = 2bqw$ . Set  $w_+ := \Pi_{q+1}^+ w$ ,  $w_- := w - w_+$ . The operator  $H_0 - \Pi_{q+1}^+ V \Pi_{q+1}^+ - 2bq$  is boundedly invertible in  $\Pi_{q+1}^+ L^2(\mathbb{R}^2)$ , its inverse is a positive operator, and by analogy with (2.10) we get  $w_+ = (H_0 - \Pi_{q+1}^+ V \Pi_{q+1}^+ - 2bq)^{-1} \Pi_{q+1}^+ V \Pi_{q+1}^- w_-$ . Further, similarly to (2.11), we find that

$$\begin{aligned} & \langle (H_0 - 2bq)w_-, w_- \rangle - \langle \Pi_{q+1}^+ V \Pi_{q+1}^- w_-, w_- \rangle \\ & - \langle \Pi_{q+1}^- V \Pi_{q+1}^+ (H_0 - \Pi_{q+1}^+ V \Pi_{q+1}^+ - 2bq)^{-1} \Pi_{q+1}^+ V \Pi_{q+1}^- w_-, w_- \rangle = 0. \end{aligned}$$

The three terms on the l.h.s. are non-positive, and hence they should vanish. As in the proof of (1.3), we easily conclude that  $w_- = 0$ , and hence  $w = 0$ .  $\square$

### 3 Proof of Theorem 2

Define the operators

$$H_0^{(m)} := -\frac{1}{\varrho} \frac{d}{d\varrho} \varrho \frac{d}{d\varrho} + \left( \frac{m}{\varrho} - b\varrho \right)^2 - b, \quad m \in \mathbb{Z},$$

self-adjoint in  $L^2(\mathbb{R}_+; \varrho d\varrho)$ , as the Friedrichs' extensions of the operators defined on  $C_0^\infty(\mathbb{R}_+)$  with  $\mathbb{R}_+ := (0, \infty)$ . Then, the operator  $H_0$  is unitarily equivalent to the orthogonal sum  $\bigoplus_{m \in \mathbb{Z}} H_0^{(m)}$  under the passage to polar coordinates  $(\varrho, \phi)$  in  $\mathbb{R}^2$ , and a subsequent decomposition into a Fourier series with respect to the angular variable  $\phi$ . For any  $m \in \mathbb{Z}$ , we have

$$\sigma(H_0^{(m)}) = \bigcup_{q=m_-}^{\infty} \{2bq\}$$

where, as usual,  $m_- := \max\{0, -m\}$  (see e.g. [1]). In contrast to the 2D Landau Hamiltonian  $H_0$  however, we have  $\dim \text{Ker}(H_0^{(m)} - 2bq) = 1$  for all  $q \geq m_-$ ,  $m \in \mathbb{Z}$ . Further, assume that  $V \in L^\infty(\mathbb{R}^2; \mathbb{R})$  and  $V$  is radially symmetric i.e.

$$V(x, y) = v\left(\sqrt{x^2 + y^2}\right), \quad (x, y) \in \mathbb{R}^2.$$

Then, the operator  $H_0 + V$  is unitarily equivalent to the orthogonal sum  $\bigoplus_{m \in \mathbb{Z}} (H_0^{(m)} + v)$ . Thus,

$$\dim \text{Ker}(H_0 + V - 2bq) = \sum_{m \in \mathbb{Z}} \dim \text{Ker}(H_0^{(m)} + v - 2bq), \quad q \in \mathbb{N}. \quad (3.1)$$

If  $\|V\|_{L^\infty(\mathbb{R}^2)} = \|v\|_{L^\infty(\mathbb{R}_+)} < b$ , for all  $m \in \mathbb{N}$ , the  $q$ -th eigenvalue of  $H_0^{(m)} + v$  that we denote by  $E_q(v; m)$ , stays in the interval  $2bq + ] - b, b[$ ; in particular, it stays simple. So, as a consequence of regular perturbation theory, see e.g. [5, 13], the eigenvalues  $(E_q(v; m))_{q \geq 0}$  are real analytic functions of the potential  $v$ . Moreover, one computes

$$\frac{\partial}{\partial t} E_q(tv; m)|_{t=0} = \int_{\mathbb{R}_+} v(\varrho) \varphi_{q,m}(\varrho)^2 \varrho d\varrho \quad (3.2)$$

where

$$\varphi_{q,m}(\varrho) := \sqrt{\frac{q!}{\pi(q+m)!} \left(\frac{b}{2}\right)^{m+1}} \varrho^m L_q^{(m)}(b\varrho^2/2) e^{-b\varrho^2/4}, \quad \varrho \in \mathbb{R}_+, \quad q \in \mathbb{N},$$

are the normalized eigenfunctions of the operator  $H_0^{(m)}$ ,  $m \in \mathbb{N}$ , and

$$L_q^{(m)}(s) := \sum_{l=0}^q \frac{(q+m)!}{(m+l)!(q-l)!} \frac{(-s)^l}{l!}, \quad s \in \mathbb{R},$$

are the generalized Laguerre polynomials.

Pick  $t \in ]-b/2, b/2[^{\mathbb{N}^*}$  and consider the potential

$$v_t(\rho) = - \sum_{j \in \mathbb{N}^*} t_{2j-1} \mathbf{1}_{[x_{2j-1}^-, x_{2j-1}^+]}(\rho) + \sum_{j \in \mathbb{N}^*} t_{2j} \mathbf{1}_{[x_{2j}^-, x_{2j}^+]}(\rho), \quad \rho \in \mathbb{R}_+, \quad (3.3)$$

where  $x_j^- := e^{-\alpha_j/2}$ ,  $x_j^+ := e^{-\beta_j/2}$ , and

$$\alpha_{2j-1} := 2^{-N(j-1/2)^2+1}, \quad \beta_{2j-1} := 2^{-Nj^2+1}, \quad \alpha_{2j} := 2^{-N(j-1/2)^2}, \quad \beta_{2j} := 2^{-Nj^2}. \quad (3.4)$$

We will choose the large integer  $N$  later on.

As, for  $j \geq 1$ , one has

$$N(j-1)^2 < N(j-1/2)^2 - 1 < N(j-1/2)^2 < Nj^2 - 1 < Nj^2 < N(j+1/2)^2 - 1,$$

we note that, for  $N$  sufficiently large, one has:

- $\|v_t\|_{L^\infty(\mathbb{R}_+)} < b$  for  $t \in ]-b/2, b/2[^{\mathbb{N}^*}$ ;
- $v_t$  vanishes identically if and only if the vector  $(t_j)_j$  vanishes identically.

For  $j \geq 1$ , define  $m_j = 2^{Nj^2} - 1$  and consider the mapping

$$\mathcal{E} : t \in ]-b/2, b/2[^{\mathbb{N}^*} \mapsto (\mathcal{E}_{2j-1}(t), \mathcal{E}_{2j}(t))_{j \geq 1} = (t_{2j} + t_{2j-1}, \tilde{E}_q(v_t; m_j))_{j \geq 1} \in ]-r, r[^{\mathbb{N}^*}$$

where

$$\tilde{E}_q(v_t; m_j) = \frac{2\pi q!}{C_j} \frac{m_j(m_j!)^2}{(q+m_j)!} \left(\frac{2}{b}\right)^{m_j+1} (E_q(v_t; m_j) - 2bq)$$

The constants  $(C_j)_j$  are going to be chosen later on.

The mapping is real analytic and we can compute its Jacobi matrix at  $t = 0$ . First, bearing in mind (3.2), (3.3), and (3.4), we easily find that

$$\begin{aligned} \partial_{t_{2j}} \mathcal{E}_{2l}(0) &= C_j^{-1} (e^{-m_l \beta_{2j}} (1 + o(1)) - e^{-m_l \alpha_{2j}} (1 + o(1))) \\ &= \begin{cases} 1 & \text{if } j = l, \\ O(e^{-2^{N|j-l|}}) & \text{if } l > j, \\ O(2^{-N|j-l|}) & \text{if } l < j, \end{cases} \\ \partial_{t_{2j+1}} \mathcal{E}_{2l}(0) &= -C_j^{-1} (e^{-m_l \beta_{2j+1}} (1 + o(1)) - e^{-m_l \alpha_{2j+1}} (1 + o(1))) \\ &= \begin{cases} -e^{-2} + O(e^{-2^{Nj}}) & \text{if } j = l, \\ O(e^{-2^{N|j-l|}}) & \text{if } l > j, \\ O(2^{-N|j-l|}) & \text{if } l < j, \end{cases} \end{aligned}$$

when one chooses  $C_j$  properly. In this formula,  $o(1)$  refers to the behavior of the function when  $N \rightarrow +\infty$  uniformly in  $l, j$ . Moreover, obviously,

$$\partial_{t_{2j-1}} \mathcal{E}_{2l-1}(0) = \partial_{t_{2j}} \mathcal{E}_{2l-1}(0) = \delta_{jl}.$$

Hence, the Jacobi matrix of the mapping  $\mathcal{E}(t)$  at  $t = 0$  can be written as  $J + E$  where  $J$  is a block diagonal matrix made of the blocks  $\begin{pmatrix} 1 & 1 \\ -e^{-2} & 1 \end{pmatrix}$  and the error matrix  $E$  is a bounded operator from  $l^\infty(\mathbb{N}^*)$  to itself with a norm bounded by  $C2^{-N}$ . So for  $N$  large enough this Jacobi matrix is invertible and, using the analytic inverse mapping theorem, we see that there exists a real analytic diffeomorphism  $\varphi$  on a ball of  $l^\infty(\mathbb{N}^*)$  centered at 0, such that

$$\mathcal{E} \circ \varphi(u) = (u_{2j} + u_{2j-1}, u_{2j} - e^{-2}u_{2j-1})_{j \geq 1} \in ]-r, r[^{\mathbb{N}^*},$$

and  $\varphi(0) = 0$ . To construct the potential  $v_t$  having the Landau level  $2bq$  as an eigenvalue with infinite multiplicity, it suffices to take  $t = \varphi(u)$  with  $u_{2j} = e^{-2}u_{2j-1} \neq 0$  for infinitely many indices  $j \in \mathbb{N}^*$ . This completes the proof of Theorem 2.

**Acknowledgements.** The authors were partially supported by the Chilean Scientific Foundation *Fondecyt* under Grants 7080135 and 1050716.

G. Raikov acknowledges also the partial support of *Núcleo Científico ICM P07-027-F* “*Mathematical Theory of Quantum and Classical Magnetic Systems*”.

## References

- [1] J. AVRON, I. HERBST, B. SIMON, *Schrödinger operators with magnetic fields. I. General interactions*, Duke Math. J. **45** (1978), 847-883.
- [2] N. FILONOV, A. PUSHNITSKI, *Spectral asymptotics of Pauli operators and orthogonal polynomials in complex domains* Comm. Math. Phys. **264** (2006), 759-772.
- [3] V. FOCK, *Bemerkung zur Quantelung des harmonischen Oszillators im Magnetfeld*, Z. Physik **47** (1928), 446-448.
- [4] V. IVRII, *Microlocal Analysis and Precise Spectral Asymptotics*, Springer monographs in Math. Springer, Berlin, 1998.
- [5] T. KATO, *Perturbation Theory for Linear Operators*, Springer, Berlin, 1980.
- [6] L. LANDAU, *Diamagnetismus der Metalle*, Z. Physik **64** (1930), 629-637.
- [7] M. MELGAARD, G. ROZENBLUM, *Eigenvalue asymptotics for weakly perturbed Dirac and Schrödinger operators with constant magnetic fields of full rank*, Comm. PDE **28** (2003), 697-736.

- [8] K. MILLER, B. SIMON, *Quantum magnetic Hamiltonians with remarkable spectral properties*, Phys. Rev. Lett. **44** (1980), 1706–1707.
- [9] M. PERSSON, *Eigenvalue asymptotics of the even-dimensional exterior Landau-Neumann Hamiltonian*, Adv. Math. Phys. **2009** (2009), Article ID 873704, 15 pp.
- [10] A. PUSHNITSKI, G. ROZENBLUM, *Eigenvalue clusters of the Landau Hamiltonian in the exterior of a compact domain*, Doc. Math. **12** (2007), 569–586.
- [11] G. D. RAIKOV, *Eigenvalue asymptotics for the Schrödinger operator with homogeneous magnetic potential and decreasing electric potential. I. Behaviour near the essential spectrum tips*, Comm. PDE **15** (1990), 407-434; Errata: Comm. PDE **18** (1993), 1977-1979.
- [12] G. D. RAIKOV, S. WARZEL, *Quasi-classical versus non-classical spectral asymptotics for magnetic Schrödinger operators with decreasing electric potentials*, Rev. Math. Phys. **14** (2002), 1051–1072.
- [13] M. REED, B. SIMON, *Methods of Modern Mathematical Physics. IV. Analysis of operators*, Academic Press, 1978.
- [14] G. ROZENBLUM, A. SOBOLEV *Discrete spectrum distribution of the Landau operator perturbed by an expanding electric potential*, to appear in: Advances in the Mathematical Sciences Spectral Theory of Differential Operators: M. Sh. Birman 80th Anniversary Collection **225** (2008), 169-190.
- [15] G. ROZENBLUM, G. TASHCHIYAN, *On the spectral properties of the perturbed Landau Hamiltonian*, Comm. Partial Differential Equations **33** (2008), 1048–1081.

FRÉDÉRIC KLOPP

Département de mathématiques et Institut Universitaire de France  
 Université de Paris Nord  
 Avenue J.Baptiste Clément  
 93430 Villetaneuse, France  
 E-mail: klopp@math.univ-paris13.fr

G. RAIKOV

Facultad de Matemáticas  
 Pontificia Universidad Católica de Chile  
 Av. Vicuña Mackenna 4860  
 Santiago de Chile  
 E-mail: graikov@mat.puc.cl